

AD-A259 398

ATION PAGE

Form Approved
OMB No. 0704-0188

②



Do not fill in this section unless you are submitting a report to the Office of Management and Budget. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Washington, DC 20540.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT TYPE AND DATES COVERED	
4. TITLE AND SUBTITLE <i>GENETICALLY EPILEPSY-PRONE RATS HAVE INCREASED BRAIN REGIONAL ACTIVITY OF AN ENZYME WHICH LIBERATES GLUTAMATE FROM NAAAG</i>				5. FUNDING NUMBERS	
6. AUTHOR(S) <i>JAMES L. MOYMHOF, RUTH E. CARTER</i>					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) WALTER REED ARMY INSTITUTE OF RESEARCH WASHINGTON, D.C. 20307-5100				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND FT. DETRICK, FREDERICK, MD. 21702-5012				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
<div style="position: relative; width: 100%; height: 100%;"> <div style="position: absolute; top: 0; left: 0; right: 0; bottom: 0; display: flex; align-items: center; justify-content: center;"> <div style="font-size: 2em; font-weight: bold; transform: rotate(-15deg);">S</div> <div style="text-align: center;"> <div style="font-size: 2em; font-weight: bold; transform: rotate(-15deg);">DTIC</div> <div style="font-size: 1.5em; font-weight: bold; transform: rotate(-15deg);">ELECTE</div> <div style="font-size: 1.5em; font-weight: bold; transform: rotate(-15deg);">JAN 2 1993</div> </div> <div style="font-size: 2em; font-weight: bold; transform: rotate(15deg);">D</div> </div> </div>					
12a. DISTRIBUTION/AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <div style="display: flex; justify-content: space-between;"> <div style="width: 30%;"> <p><i>SEE ATTACH.</i></p> </div> <div style="width: 70%;"> <p>N-Acetylated-α-linked acidic dipeptidase (NAALADase), is a membrane-bound peptidase which hydrolyzes the endogenous neuropeptide N-acetyl-aspartyl-glutamate (NAAg) to N-acetyl-aspartate (NAA) and the excitatory amino acid, glutamate (GLU). Although there is evidence that NAAG might be a neurotransmitter, this dipeptide could also function as a precursor form of GLU, which is liberated by the dipeptidase. We found that the activity of this NAAG hydrolyzing enzyme in genetically epilepsy-prone rats was 11-26% greater than control in brain regions, including the amygdala, hippocampus and cerebellum, as well as the pyriform, entorhinal and frontal cortices. This is consistent with possible increased availability of GLU in certain CNS synapses in these rats, which are reported to have increased susceptibility to audiogenically, electrically and chemically induced convulsions.</p> </div> </div>					
14. SUBJECT TERMS <i>GENETICS, HIPPOCAMPUS, EPILEPSY, NAAAG</i>				15. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT		18. SECURITY CLASSIFICATION OF THIS PAGE		19. SECURITY CLASSIFICATION OF ABSTRACT	
				20. LIMITATION OF ABSTRACT	

BRES 25401

Sof

93-00636



Genetically epilepsy-prone rats have increased brain regional activity of an enzyme which liberates glutamate from *N*-acetyl-aspartyl-glutamate¹

James L. Meyerhoff^a, Ruth E. Carter^b, Debra L. Yourick^a, Barbara Stauch Slusher^c
and Joseph T. Coyle^d

^a Department of Medical Neurosciences, Division of Neuropsychiatry, Walter Reed Army Institute of Research, Washington, DC 20307 (USA),

^b Department of Pharmacology and Molecular Sciences, Johns Hopkins University School of Medicine, Baltimore, MD 21205 (USA);

^c Department of CNS Pharmacology, ICI Americas, Wilmington, DE 19897 (USA) and ^d Department of Psychiatry, Harvard Medical School, Belmont, MA 02178 (USA)

(Accepted 21 July 1992)

Key words: Genetics; Hippocampus; *N*-Acetylasparylglutamate Epilepsy; Glutamate; *N*-Acetylated- α -linked acidic dipeptidase; Enzymatic; Hydrolysis; NAAG; Aspartate

N-Acetylated- α -linked acidic dipeptidase (NAALADase) is a membrane-bound peptidase which hydrolyzes the endogenous neuropeptide *N*-acetylasparylglutamate (NAAG) to *N*-acetylaspartate (NAA) and the excitatory amino acid, glutamate (Glu). Although there is evidence that NAAG might be a neurotransmitter, this dipeptide could also function as a precursor form of Glu, which is liberated by the dipeptidase. We found that the activity of this NAAG hydrolyzing enzyme in genetically epilepsy-prone rats was 11–26% greater than control in brain regions, including the amygdala, hippocampus and cerebellum, as well as the pyriform, entorhinal and frontal cortices. This is consistent with possible increased availability of Glu in certain CNS synapses in these rats, which are reported to have increased susceptibility to audiogenically, electrically and chemically induced convulsions.

Injection of the dipeptide *N*-acetylasparylglutamate (NAAG) into rat hippocampus causes prolonged seizure activity, similar to that seen following infusion of quisqualic acid, a glutamate agonist²³. NAAG is a dipeptide found in the CNS in high concentrations^{4,10,14}. NAAG is hydrolyzed to glutamate (Glu) and *N*-acetylaspartate (NAA) by *N*-acetylated- α -linked acidic dipeptidase (NAALADase), a chloride-dependent peptidase, enriched in synaptic membranes¹⁷. Despite evidence supporting the view of NAAG as a putative neurotransmitter, the identification, characterization and distribution of NAALADase suggest an alternative hypothesis: NAAG may function as a storage form of Glu, which could be liberated by this peptidase and released into the synaptic cleft^{2,12,19,20}. We previously reported that non-kindled convulsions are as effective

as kindled seizures in decreasing the activity of this enzyme in the amygdala, pyriform cortex, frontal cortex and entorhinal cortex¹². This indicates that the decreases seen in the kindled group should be attributed to the aftereffects of convulsive activity rather than to the kindling process. Such decreases could possibly serve as a homeostatic mechanism, limiting CNS excitability by restricting the availability of Glu at synapses where NAAG might be the precursor. Extending this hypothesis, we asked whether the activity of this dipeptidase might be elevated in conditions associated with increased excitability, such as genetic epilepsy. In a pilot investigation of this question, we found elevated NAALADase activity in homogenates from brain regions of adult progeny inbred from severe-seizure genetically epilepsy-prone rats (GEPR-9s), compared to

Correspondence to: J.L. Meyerhoff, Department of Medical Neurosciences, Walter Reed Army Institute of Research, Washington, DC 20307, USA.

¹ Research was conducted in compliance with the Animal Welfare Act, and other Federal statutes and regulations relating to animals and experiments involving animals and adheres to principles stated in the *Guide for the Care and Use of Laboratory Animals*, NIH publication 86-23, 1985 edition. The views of the author(s) do not purport to reflect the position of the Department of the Army or the Department of Defense, (para 4-3, AR 360-5).

98 1 11 004

controls (unpublished data). The present experiment is a replication of the pilot study with two refinements: (1) to ensure positive phenotype in 100% of the GEPR-9s delivered, rats were screened by the supplier for tonic extensor convulsive responses to audiogenic challenge; (2) NAALADase assays were carried out in membrane preparations, which yield higher activity of this membrane-bound enzyme than do homogenates^{2,19}.

Adult male GEPR-9^{5,15} and control¹⁵ rats (12–18 weeks of age, 350–500 g, obtained from Dr. John Dailey, Department of Basic Sciences, University of Illinois College of Medicine, Peoria, IL 61656) were maintained in a sound-attenuating animal enclosure. The development, testing and screening of these colonies is detailed by Reigel et al.¹⁵. GEPR-9s were derived from Sprague–Dawley rats, inbred and screened to select for reliable tonic convulsive responses to audiogenic challenge. Ninety-five percent of audiogenically screened GEPR-9 progeny meet the criteria for susceptibility to severe audiogenic seizures^{5,15}, and only these rats are shipped as screened GEPR-9s. Control rats were derived from Sprague–Dawley rats, inbred and screened to select for reliable resistance to audiogenic challenge¹⁵, and all animals shipped were seizure-free on three separate audiogenic challenges (Dr. John Dailey, personal communication). After shipment to the Walter Reed Army Institute of Research, the rats were individually housed for over 80 days in a temperature-controlled room with a 12 h/12 h light/dark cycle, and food and water were freely available.

Rats were killed by decapitation, alternating GEPR-9s and controls. The brains were removed and immediately dissected on ice. The dissection was derived from a method previously published¹¹. The regional samples were weighed, rapidly frozen on dry ice and stored at -70°C until assayed. Samples were identified as to brain region, but treatment groups were coded, and samples were assayed without identification of treatment group.

To assay for NAALADase activity, tissue samples were homogenized (six strokes, 600 r.p.m.) in ice-cold 50 mM Tris-HCl buffer (pH 7.4 at 37°C) at a dilution of 1:100 (wet weight of tissue (g)/volume of buffer (ml)). An aliquot of the homogenate was sonicated for 5 s to disrupt cellular membranes and centrifuged for 20 min at 4°C at 35,000 g. The entire pellet was then resuspended in 50 mM Tris-HCl + 0.5% Triton X-100 by sonication. The rest of the radioenzymatic assay¹⁷ was performed as modified by Slusher et al.¹⁹. Due to the ionic sensitivity of enzyme activity we substituted triple-distilled, deionized water (purchased from J.T. Baker, Inc., Phillipsburg, NJ), for double-distilled, deionized, tap water in all assay reagents. In brief, the assay measured the amount of [^3H]Glu liberated from [^3H]NAAG during a 15 min incubation at 37°C , in 50 mM Tris-HCl buffer, pH 7.4, with Triton X-100 in a final concentration of 0.05%. Substrate and product were separated (95% resolution) via anion-exchange liquid chromatography. Samples were assayed in triplicate. NAALADase activity was calculated based on picomoles of [^3H]Glu produced from [^3H]NAAG per mg protein per min. The data were decoded and analyzed by a one-way analysis of variance.

As shown in Table I, NAALADase activity was greater in GEPR-9s compared to controls in membrane preparations from each of the brain regions examined. Activity was higher in the pyriform (19%), entorhinal (24%) and frontal (26%) cortices, as well as in the hippocampus (18%), amygdala (17%) and cerebellum (11%). Several of these regions are involved in limbic seizures. It is unlikely that spontaneous seizures affected these results, as they rarely occur in the audiogenically seizure-sensitive GEPR-9¹⁵. Moreover, since convulsions decrease NAALADase activity¹², if they did occur, they would lead to an underestimation of the increase we find in GEPRs. GEPR-9s and controls in this study had been screened at the University of Illinois on three occasions between 45–65 days of age for the presence (GEPR-9) or absence (control) of

TABLE I
NAALADase activity in GEPR-9 vs control rat brain membranes

Region	Control (pmol per mg protein per min \pm SEM)	GEPR-9	F	P	N	Increase (%)
Amygdala	25.6 \pm 1.8	30.0 \pm 1.4	6.8	< 0.05	10	17
Pyriform Cortex	25.1 \pm 1.3	23.8 \pm 0.9	19.0	< 0.002	10	19
Entorhinal Cortex	22.0 \pm 0.9	27.3 \pm 1.3	44.3	< 0.001	10	24
Frontal Cortex	18.0 \pm 1.1	22.7 \pm 0.9	47.0	< 0.001	10	26
Hippocampus	23.8 \pm 1.0	28.0 \pm 1.0	41.8	< 0.001	10	18
Cerebellum	30.4 \pm 1.1	33.8 \pm 0.8	33.4	< 0.001	10	11

tonic convulsive responses to audiogenic challenge¹⁵. But it is unlikely that the convulsive responses of GEPR-9s during screening would have affected the results, because almost 3 months elapsed between the audiogenic testing and the sacrifice of the animals. Furthermore, we statistically compared the percent differences from the present study with those seen in the pilot study done in regional brain homogenates from unscreened GEPR-9s (which never had convulsions) and controls. It may be inferred that only 5% of the unscreened GEPR progeny would have failed to meet the criteria for GEPR-9 phenotype, had they been tested¹⁵. In the pilot study, NAAG-hydrolyzing activity was significantly elevated in the pyriform (21%), entorhinal (27%) and frontal (19%), cortices, as well as in the hippocampus (29%), amygdala (8%) and cerebellum (15%). These percent elevations, in homogenates from unscreened GEPRs in the pilot study, are not distinguishable by ANOVA from the percent elevations seen in membrane preparations from screened GEPR-9s in the present study (Table I).

GEPR-9s have been found to be hypothyroid, with abnormally low serum levels of thyroxine up to day 22 and low levels of triiodothyronine throughout life¹³. Abnormalities of the auditory system have also been reported in GEPRs, but Reigel, et al.¹⁶, point out that seizure susceptibility is associated with neural structures outside that system, as well. Moreover, GEPRs also have increased susceptibility to electrically and chemically induced seizures, and the latter susceptibility appears developmentally prior to audiogenic susceptibility¹⁶. Numerous studies of neurochemical abnormalities in the GEPR-9 have described deficiencies in central noradrenergic^{6,22} and serotonergic⁹ activity. In recent years, however, epilepsy research has increasingly focussed on the possible role of putative excitatory amino acid neurotransmitters such as Glu⁸ and aspartate (ASP)⁷. NAAG has been proposed to be a neurotransmitter/neuromodulator at a subpopulation of Glu/ASP synapses. A significant number of studies, employing neurochemical, immunocytochemical, stimulation and lesioning techniques, as well as demonstrations of calcium-dependent, stimulated release of NAAG have supported this hypothesis (see review by Coyle et al.³). Of particular relevance to epilepsy is the description of NAAG-immunoreactive neurons in the entorhinal cortex¹, a region in which NAAG levels were increased by kindling¹¹.

NAALADase appears to be the primary mechanism for catabolism of NAAG in brain¹⁹. It has been shown to hydrolyze NAAG into NAA and Glu both *in vitro*¹⁷ and *in vivo*^{2,21}. Activity is weakly inhibited by the reaction products Glu and NAA^{17,19}. This peptidase

has been solubilized from synaptosomal membranes and purified to apparent homogeneity¹⁹. The purified enzyme migrates as a single band on an SDS gel with an apparent MW of 94 kDa, and displays a remarkably high apparent affinity for its putative substrate, NAAG, with a K_m of 140 nM¹⁹. The enzyme has been immunocytochemically localized in the CNS²⁰, where it is selectively enriched in many structures known to contain NAAG. Developmentally, the activity of this peptidase in the cerebral cortex of normal rats reaches 80% of maximum at 21 days of age², a developmental milestone which coincides with the period of synaptogenesis in normal rats, as well as the age at which seizure incidence in response to audiogenic challenge first reaches 100% in GEPR-9s¹⁶. Although some evidence suggests that NAAG might be a neurotransmitter, more recently it has been proposed that the function of NAALADase *in vivo* may be to liberate a neuroactive fragment, such as Glu from its dipeptide precursor^{2,12,19}. Therefore, assuming access to substrate, the elevated activity of this enzyme seen in the GEPR-9 might increase availability of NAA and Glu in certain synapses of the brain. While the increase in Glu is expected to be excitatory, there is no indication that NAA would have any effect on neurotransmission².

Further experiments are required to fully assess the implications of the present study. We anticipate conducting studies to determine the effect of thyroidectomy on the ontogeny of NAAG hydrolyzing activity, as well as an expanded regional analysis of the activity of this peptidase in strains of seizure-prone and seizure-resistant rats and mice to determine whether abnormally high activity of this enzyme is always associated with genetically determined seizure susceptibility¹⁸.

The authors are grateful to Dr. Frank Tortella, Dr. G. Jean Kant and Dr. Michael Koenig for critical reading of the manuscript. We also thank SGT Edward F. Knight, Ron Repasi, Chris Hogan, Thuan Dang, Rob Greenwald and Kerri-Lynn Primmer for technical assistance.

- 1 Anderson, K.A., Monaghan, D.T., Cangro, C.B., Nambodiri, M.A.A., Neale, J.A. and Cotman, C.W., Localization of N-acetylaspartylglutamate-like immunoreactivity in selected areas of the rat brain, *Neurosci. Lett.*, 72 (1986) 14-20.
- 2 Blakely, R.D., Robinson, M.B., Thompson, R.C. and Coyle, J.T., Hydrolysis of the brain dipeptide N-acetyl-L-aspartyl-L-glutamate: Subcellular and regional distribution, ontogeny, and the effect of lesion on N-acetylated alpha linked acidic dipeptidase activity, *J. Neurochem.*, 50 (1988) 1200-1209.
- 3 Coyle, J.T., Stauch-Slusker, B., Tsai, G., Rothstein, J., Meyerhoff, J.L., Simmons, M. and Blakely, R.D., N-acetylaspartyl glutamate. In B.S. Meldrum, F. Moroni, R.P. Simon, and J.H. Woods (Eds.), *Excitatory Amino Acids*. Raven Press, New York, 1991, pp. 69-77.
- 4 Curatolo, A., D'Archangelo, P., Lino, A. and Brancati, A., Distribution of N-acetylaspartic and N-acetylaspartylglutamic acids in nervous tissue, *J. Neurochem.*, 12 (1965) 339-342.

- 5 Dailey, J.W. and Jobe, P.C., Anticonvulsant drugs and the genetically epilepsy-prone rat, *Fed. Proc.*, 44 (1985) 2640-2644.
- 6 Dailey, J.W. and Jobe, P.C., Indices of noradrenergic function in the central nervous system of seizure-naive genetically epilepsy-prone rats, *Epilepsia*, 27 (1986) 665-670.
- 7 Flavin, H.J., Wieraszko, A. and Seyfried, T.N., Enhanced aspartate release from hippocampal slices of epileptic (E1) mice, *J. Neurochem.*, 56 (1991) 1007-1011.
- 8 Geula, C., Jarvie, P.A., Logan, T.C. and Slevin, J.T., Long-term enhancement of K⁺-evoked release of L-glutamate in entorhinal kindled rats, *Brain Res.* 442 (1988) 368-372.
- 9 Jobe, P.C. and Laird, H.E., Neurotransmitter abnormalities as determinants of seizure susceptibility and intensity in the genetic models of epilepsy, *Biochem. Pharmacol.*, 30 (1981) 3137-3144.
- 10 Koller, K.J., Zaczek, R. and Coyle J.T., N-acetylaspartylglutamate: regional levels in rat brain and the effects of brain lesions as determined by a new HPLC method, *J. Neurochem.*, 43 (1984) 1136-1142.
- 11 Meyerhoff, J.L., Koller K.J., Walczak, D.D. and Coyle, J.T., Brain regional levels of N-acetylaspartylglutamate (NAAG): The effect of kindled seizures, *Brain Res.*, 346 (1985) 392-396.
- 12 Meyerhoff, J.L., Robinson, M.B., Bixler, M.A., Richards, S.S. and Coyle, J.T., Seizures decrease regional enzymatic hydrolysis of N-acetylaspartylglutamate in rat brain, *Brain Res.*, 505 (1989) 130-134.
- 13 Mills, S.A. and Savage, D.D., Evidence of hypothyroidism in the genetically epilepsy-prone rat, *Epilepsy Res.*, 2 (1988) 102-110.
- 14 Reichert K.L. and Fonnum F., Subcellular localization of N-acetylaspartylglutamate, N-acetylglutamate and glutathione in brain, *J. Neurochem.*, 16 (1969) 1409-1416.
- 15 Reigel, C.E., Dailey, J.W. and Jobe, P.C., The genetically epilepsy-prone rat: an overview of seizure-prone characteristics and responsiveness to anticonvulsant drugs, *Life Sci.*, 39 (1986) 763-774.
- 16 Reigel, C.E., Jobe, P.C., Dailey, J.W. and Savage, D.D., Ontogeny of sound-induced seizures in the genetically epilepsy-prone rat, *Epilepsy Res.*, 4 (1989) 63-71.
- 17 Robinson, M.B., Blakely, R.D., Couto, R. and Coyle, J.T., Hydrolysis of the brain dipeptide N-acetyl-L-aspartyl-L-glutamate: Identification and characterization of a novel N-acetylated-alpha-linked acidic dipeptidase activity from rat brain, *J. Biol. Chem.*, 262 (1987) 14498-14506.
- 18 Seyfried, T.N., Glaser, G.H., Yu, R.K. and Palayoor, S.T., Inherited convulsive disorders in mice, *Adv. Neurol.*, 44 (1986) 115-133.
- 19 Slusher, B.S., Robinson, M.B., Tsai, G., Simmons, M.L., Richards, S.S. and Coyle, J.T., Rat brain N-acetylated α -linked acidic dipeptidase activity, *J. Biol. Chem.*, 265 (1990) 21297-21301.
- 20 Slusher, B.S., Tsai, G., Yoo, G. and Coyle, J.T., Immunocytochemical localization of the N-acetylaspartylglutamate (NAAG) hydrolyzing enzyme N-acetylated α -linked acidic dipeptidase (NAALADase), *J. Comp. Neurol.*, 315 (1992) 217-229.
- 21 Stauch, B.S., Robinson, M.B., Forloni, G., Tsai, G. and Coyle, J.T., The effects of N-acetylated alpha-linked acidic dipeptidase (NAALADase) inhibitors on [³H]NAAG catabolism in vivo, *Neurosci. Lett.*, 100 (1989) 295-300.
- 22 Yourick, D.L., La Placa, M.C. and Meyerhoff, J.L., Norepinephrine-stimulated phosphatidylinositol metabolism in genetically epilepsy-prone and kindled rats, *Brain Res.*, 551 (1991) 315-318.
- 23 Zaczek, R., Koller, K., Cotter, R., Heller, D. and Coyle, J.T., N-acetylaspartylglutamate: An endogenous peptide with high affinity for a brain 'glutamate' receptor, *Proc. Natl. Acad. Sci. USA* 80 (1983) 1116-1119.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	20

DTIC QUALITY INSPECTED 1